

**AMENDMENTS TO THE SPECIFICATION:**

*Please amend the paragraph at page 8, line 5, as follows:*

**[Cationically ~~Photopolymerizable~~ Polymerizable Compound (B)]**


The cationically polymerizable compound (B) in the invention is a compound having at least one cationically polymerizable functional group in one molecule, and specifically a material selected from a compound containing an epoxy group, a compound containing an oxetanyl group, an oxolane compound, a cyclic acetal compound, a cyclic lactone compound, a thiirane compound, a thietane compound, a spiroorthoester compound, a vinyl ether compound, an ethylenically unsaturated compound, a cyclic ether compound, a cyclic thioether compound and a vinyl compound. Preferably used is a material containing an epoxy group or an oxetanyl group as a functional group.

*Please amend the paragraph at page 14, line 5, as follows:*

**[Cyclic polyether compound (C)]**

As the cyclic ~~polyoxyethylene~~ polyether compound (C) according to the invention, mention may be made of the compound represented by the following formula (6).

Formula (6):



*Please amend the paragraph at page 16, line 17, as follows:*

In the formula (7) or (8), R<sub>2</sub> is an ~~alkylene~~ alkyl group such as a methyl group, an ethyl group or a propyl group. R<sub>3</sub> is a linear or branched alkylene group containing 1 to 20 carbon atoms such as a methylene group, an ethylene group, a propylene group or a butylene group; a linear or branched poly(alkyleneoxy) group containing 1 to 120 carbon atoms such as a poly(ethyleneoxy) group or a poly(propyleneoxy) group; a linear or branched unsaturated hydrocarbon group such as a propenylene group, a methylpropenylene group or a butenylene group; a carbonyl group; an alkylene group having a carbonyl group; an alkylene group having a carbamoyl group in the molecular chain; or a phenyl group. Further, R<sub>4</sub> is an ~~alkylene~~ alkyl group such as a methyl group, an ethyl group or a propyl group; a glycidyl ether group; primary amine; a thiol group; a vinyl group; or an isocyanate group. Materials in which the hydrogen atoms of the carbon-hydrogen bonds in these coupling agents are partially or completely substituted with fluorine atoms can also be used. Preferably, materials in which the hydrogen atom in a methylene group or a methyl group is partially or completely substituted with a fluorine atom are used.

*Please amend Table 1 at page 26, as follows:*

[illegible]